

Epidemiological and clinical insights from SARS-CoV-2 RT-PCR crossing threshold values, France, January to November 2020

Samuel Alizon^{1*}, Christian Selinger^{2*}, Mircea T Sofonea^{3*}, Stéphanie Haim-Boukoba³, Jean-Marc Giannoli⁴, Laetitia Ninove⁵, Sylvie Pillet^{6,7}, Vincent Thibault⁸, Alexis de Rougemont^{9,10}, Camille Tumiotto¹¹, Morgane Solis¹², Robin Stephan¹³, Céline Bressollette-Bodin¹⁴, Maud Salmona¹⁵, Anne-Sophie L'Honneur¹⁶, Sylvie Behillil¹⁷, Caroline Lefeuvre^{18,19}, Julia Dina²⁰, Sébastien Hantz^{21,22}, Cédric Hartard²³, David Veyer²⁴, Héloïse M Delagrèverie²⁵, Slim Fourati²⁶, Benoît Visseaux²⁷, Cécile Henquell²⁸, Bruno Lina²⁹, Vincent Foulongne³⁰, Sonia Burrel³¹, on behalf of the SFM COVID-19 study group³²

1. MIVEGEC, CNRS, IRD, Université de Montpellier, France
2. Center for Interdisciplinary Research in Biology (CIRB), Collège de France, CNRS, INSERM, Université PSL, Paris, France
3. Laboratoire CERBA, Saint-Ouen-L'Aumône, France
4. BIOGROUP, Scientific Direction, Lyon, France
5. Unité des Virus Émergents (UVE: Aix-Marseille Univ-IRD 190-Inserm 1207-IHU Méditerranée Infection), Marseille, France
6. Laboratoire des agents infectieux et d'hygiène, CHU de Saint-Etienne, France
7. CIRI, Centre International de Recherche en Infectiologie, GIMAP team, University of Lyon, University of Saint-Etienne, INSERM, U1111, CNRS UMR5308, ENS de Lyon, UCBL, Lyon, France
8. Laboratoire de Virologie, CHU Rennes, Rennes, France
9. Laboratory of Virology-Serology, University Hospital of Dijon Bourgogne, Dijon, France
10. UMR PAM A 02.102 Procédés Alimentaires et Microbiologiques, Université de Bourgogne Franche-Comté/AgroSup Dijon, Dijon, France
11. University of Bordeaux, CNRS-UMR 5234, CHU Bordeaux, Virology Department, Bordeaux, France
12. CHU de Strasbourg, Laboratoire de Virologie, Strasbourg, France, Université de Strasbourg, INSERM, IRM UMR_S 1109, Strasbourg, France
13. Laboratoire de Microbiologie, CHU Nîmes, Nîmes, France
14. CHU Nantes, Nantes Université, Service de Virologie, Nantes, France
15. 15 Laboratoire de Virologie, Hôpital Saint Louis, APHP, INSERM U976, équipe INSIGHT, Université de Paris, Paris, France
16. Assistance Publique-Hôpitaux de Paris, Hôpital Cochin, Service de Virologie, Paris, France
17. 17 National Reference Center for Respiratory Viruses, Molecular Genetics of RNA Viruses, UMR 3569 CNRS, University of Paris, Institut Pasteur, Paris, France
18. Département de Biologie des Agents Infectieux, Laboratoire de Virologie, CHU d'Angers, Angers, France
19. Laboratoire HIFIH, UPRES EA 3859, Université d'Angers, Angers, France
20. Laboratoire de Virologie, CHU de Caen, UNICAEN, INSERM U1311 DYNAMICURE, Université de Caen Normandie, Caen, France
21. CHU Limoges, Laboratoire de Bactériologie-Virologie-Hygiène, Limoges, France
22. RESINFIT, U 1092, University of Limoges, Limoges, France
23. Laboratoire de Virologie, CHRU de Nancy Brabois, Vandoeuvre-lès-Nancy, France; Université de Lorraine, CNRS, LCPME, Nancy, France
24. Laboratoire de Virologie, Service de Microbiologie, hôpital européen Georges Pompidou, Assistance Publique-Hôpitaux de Paris et Unité de Génomique Fonctionnelle des Tumeurs Solides, Centre de Recherche des Cordeliers, INSERM, Université Paris, Paris, France.
25. AP-HP, Hôpital Avicenne, Laboratoire de microbiologie clinique, Bobigny, France
26. Henri Mondor Hospital, virology department, Créteil, France
27. Université de Paris, Inserm, UMR 1137 IAME et Laboratoire de Virologie, Hôpital Bichat Claude Bernard, AP-HP, Paris, France
28. Service de Virologie médicale, 3IHP, CHU Clermont-Ferrand, Clermont-Ferrand, France
29. CNR des virus des infections respiratoires (dont la Grippe), Institut des Agents Infectieux, Hôpital de la Croix Rousse, HCL, Lyon, France
30. Pathogenesis and control of chronic and emerging infections, Université de Montpellier, UMR 1058, CHU de Montpellier, Inserm, Université des Antilles, Montpellier, France
31. Sorbonne Université, INSERM U1136, Institut Pierre Louis d'Epidémiologie et de Santé Publique (IPLESP), AP-HP, Hôpital Pitié-Salpêtrière, Service de Virologie, Paris, France
32. French Society of Microbiology (SFM), <https://www.sfm-microbiologie.org>

* These authors contributed equally to this work.

Correspondence: Samuel Alizon (samuel.alizon@cnrs.fr)

Citation style for this article:

Alizon Samuel, Selinger Christian, Sofonea Mircea T, Haim-Boukoba Stéphanie, Giannoli Jean-Marc, Ninove Laetitia, Pillet Sylvie, Thibault Vincent, de Rougemont Alexis, Tumiotto Camille, Solis Morgane, Stephan Robin, Bressollette-Bodin Céline, Salmona Maud, L'Honneur Anne-Sophie, Behillil Sylvie, Lefeuvre Caroline, Dina Julia, Hantz Sébastien, Hartard Cédric, Veyer David, Delagrèverie Héloïse M, Fourati Slim, Visseaux Benoît, Henquell Cécile, Lina Bruno, Foulongne Vincent, Burrel Sonia, on behalf of the SFM COVID-19 study group. Epidemiological and clinical insights from SARS-CoV-2 RT-PCR crossing threshold values, France, January to November 2020. *Euro Surveill*. 2022;27(6):pii=2100406. <https://doi.org/10.2807/1560-7917.ES.2022.27.6.2100406>

Article submitted on 31 Mar 2021 / accepted on 15 Oct 2021 / published on 10 Feb 2022

Background: The COVID-19 pandemic has led to an unprecedented daily use of RT-PCR tests. These tests are interpreted qualitatively for diagnosis, and the relevance of the test result intensity, i.e. the number of quantification cycles (Cq), is debated because of strong potential biases. **Aim:** We explored the possibility to use Cq values from SARS-CoV-2 screening tests to better understand the spread of an epidemic and to better understand the biology of the infection. **Methods:** We used linear regression models to analyse a large database of 793,479 Cq values from tests performed on more than 2 million samples between 21 January and 30 November 2020, i.e. the first two pandemic waves. We performed time series analysis

using autoregressive integrated moving average (ARIMA) models to estimate whether Cq data information improves short-term predictions of epidemiological dynamics. **Results:** Although we found that the Cq values varied depending on the testing laboratory or the assay used, we detected strong significant trends associated with patient age, number of days after symptoms onset or the state of the epidemic (the temporal reproduction number) at the time of the test. Furthermore, knowing the quartiles of the Cq distribution greatly reduced the error in predicting the temporal reproduction number of the COVID-19 epidemic. **Conclusion:** Our results suggest that Cq values of screening tests performed in the general population

TABLE

Main factors affecting Cq values of SARS-CoV-2 RT-PCR in the multivariate linear model, France, January –November 2020 (n = 793,479)

Factor	Value	Coefficient	2.5% CI	97.5% CI
Intercept		19.1	12.9	25.4
Assay	PerkinElmer		Reference	
	Genefinder	12.1	10.3	13.9
Laboratory	Lab_1		Reference	
	Lab_122	5.42	3.79	7.05
	Lab_96	-4.8	-6.71	-2.90
Result	Positive		Reference	
	Weakly positive	11.3	11.1	11.5
	Negative	16.9	16.6	17.2
Days post symptom onset	Less than 4		Reference	
	4 to 7	2.76	2.66	2.86
	8 to 14	4.90	4.73	5.08
	More than 14	5.73	5.43	6.03
Sample	Nasopharyngeal		Reference	
	Other	-1.81	-2.49	-1.14
Age	Per 20 years older	-0.541	-0.585	-0.497
Target gene	N		Reference	
	ORF1	1.03	0.949	1.12
	S	1.19	0.948	1.43
Date	Per 71 days later	-0.797	-0.903	-0.691

CI: confidence interval; Cq: quantification cycle; ORF: open reading frame; SARS-CoV-2: severe acute respiratory syndrome coronavirus 2. We only list factors with significant effects with a 10⁻³ p value criterion. Coefficients reflect differences in Cq. For qualitative factors, the reference value is shown. See the Supplement for details about the model and the scaling of the quantitative variables.

generate testable hypotheses and help improve short-term predictions for epidemic surveillance.

Introduction

Molecular testing is a key component of screening policies to control the spread of infectious diseases, and the coronavirus disease (COVID-19) pandemic has led to an unprecedented testing rate using reverse transcription PCR (RT-PCR) assays [1]. In clinical and public health practice, RT-PCR results are qualitative for viral respiratory disease diagnostics, with reports such as ‘positive’, ‘negative’, ‘uninterpretable’ and, sometimes, ‘weakly positive’. These are based on the cycles threshold, also referred to as crossing point or crossing threshold (here denoted quantification cycles (Cq)), which corresponds to the number of PCR amplification cycles required for the fluorescent signal to rise above a positive threshold. In theory, the more abundant the genetic target in the sample, the fewer the amplification cycles required to detect it. This is why numerous studies on severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) rely on Cq values to assess transmissibility [2] or study infection kinetics [3]. However, many practical and biological limitations make Cq values a poor reflector of virus load [4].

Few studies analyse Cq values at a population level. One explanation is that these are known to suffer from several, potentially strong, biases. Firstly, sample

type and sampling quality directly affect the amount of genetic material available. Secondly, the choice of RT-PCR assay matters. Even the quality of the reagents may have a considerable effect on the number of amplification cycles required to achieve the same level of fluorescence for the same amount of target genetic material. Combining data from different laboratories helps control for these sources of variation in statistical analyses. Furthermore, the larger the dataset, the more we can detect small statistical trends even after having controlled for non-informative variables.

Here, we present a cross-sectional analysis of SARS-CoV-2 RT-PCR tests performed on samples from 2,220,212 individuals in France during the COVID-19 pandemic between 21 January and 30 November 2020 (Supplementary Figure S2 shows the daily number of tests and the number of tests per French department). Our aim was to determine firstly, if this analysis at the population level can identify cofactors of interests (e.g. age, sex, type of sample) and, therefore, new virological or immunological hypotheses and, secondly, if it can improve our ability to anticipate epidemic trends.


Methods

We studied SARS-CoV-2 RT-PCR tests performed in 2020 in France on samples from individuals aged between 1 and 89 years. The national French database for SARS-CoV-2 RT-PCR tests (SI-DEP) collects qualitative results



FIGURE 1

Correlations between key factors and observed Cq variations, SARS-CoV-2 RT-PCR tests, France, January–November 2020 (n = 793,479)



Cq: quantification cycle; F: female; M: male; ORF: open reading frame; SARS-CoV-2: severe acute respiratory syndrome coronavirus 2.


For panels A, B and D, the violin plots indicate the distributions and the box plots show the 0.025, 0.25, 0.5, 0.75 and 0.975 quantiles. Panels C, E and F are obtained with a 'loess' smoothing model and the grey area shows the 95% confidence interval from the underlying model. The Cq values shown are not the raw values but those estimated using a multiparametric linear model to correct for biases (see Supplementary Methods for details regarding the computation of the estimated values).

but Cq values are not reported. In order to focus on Cq values, this study relied on the French Society of Microbiology (SFM) network of hospital-based and private laboratories. Nationwide, databases from 21 laboratories, listed in the Supplement, were included on a voluntary basis. The geographical coverage of the tests is shown in Supplementary Figure S2B. The context in which these tests were performed varied over time. Until at least April 2020, the testing capacity was limited in the country and the majority of tests were performed on symptomatic individuals, especially in hospital settings. After May 2020, testing was more accessible and data then included screening tests performed in the general population. This change in testing context coincided with a shift in terms of screening facilities, with the majority of the tests being performed

in hospital virology departments until April 2020 and in private laboratories after that. However, we do not expect that this shift led to a change in testing practice across French regions. We did not include tests for which key variables such as patient age, patient sex, laboratory geographical department, qualitative result or RT-PCR assay used were unknown. Note that one test could provide more than one Cq value if it contained probes targeting multiple viral genes. According to the national guidelines [5], it is recommended to focus on the most sensitive target to categorise levels of viral excretion. After removing the 388 Cq values that we deemed marginal and potentially unrealistic because they were smaller than 10 or larger than 45, the 95% confidence interval (CI) of the remaining values was 16.89–35.56 (details on the characteristics of the

FIGURE 2

Predicting temporal reproduction number from time series related to SARS-CoV-2 RT-PCR tests, France, January–November 2020 (n = 330,611)



Cq: quantification cycle; Rt: temporal reproduction number; SARS-CoV-2: severe acute respiratory syndrome coronavirus 2.

Panels A–D show the 7-day rolling averages of the time series of the ratio of positive tests (in purple), the median (in green) and skewness (in cyan) of the daily Cq residual distribution and Rt (in black). Panels E–F show the error made by a prediction using only Rt data (red dots) and the potential improvement made by including exogeneous data.

whole dataset and its variables of interest are included in Supplementary Table S1). The median and upper bound of the 95% CI were unaffected by the removal of these values and the lower bound increased marginally from 16.87 to 16.89. Overall, we were left with 793,479 tests from the same number of individuals, i.e. 35.7% of the whole database. The whole database contains tests with both negative and positive clinical results. However, we only kept tests with a Cq value and the former were less represented in the final database since samples that test negative usually do not have any reported Cq value (laboratories rarely record Cq values greater than 40).

We used a linear regression model to explore how Cq values can be explained by the following variables: patient age and sex, the number of days since the onset of symptoms (if known), the clinical sampling site (if known), the sampling facility (if known), the RT-PCR assay used, the target gene, the test's qualitative result, the sampling date, the temporal reproduction number (R_t) of the epidemic on the sampling date, and a control variable. The latter corresponds to the last digit of the patient anonymity number and is expected to be independent of the Cq value. Therefore, the lowest p value associated with the control variable, which we expect to be pure noise, can be used to set the significance threshold for the other variables. We also included in the model an interaction term between sampling date and R_t . For this analysis, we excluded Cq values from internal controls. Univariate analyses are extremely sensitive to heterogeneity in the dataset. For instance, the age distribution from patients sampled in aged care homes is different from that in city screening facilities, and analysing the 'sampling facility' factor alone could yield misleading results. This is why the analysis used here was multivariate and considered all the factors listed above simultaneously. In particular, it allowed us to control for variations in the way the data were collected, e.g. the intensity or the context of the sampling.

To control for the consistency of the results for some of the factors, especially those related to the infection (e.g. the number of days since symptoms onset), we also performed the analyses only on the tests that were reported as 'positive' or 'weakly positive' (i.e. we ignored the tests labelled as negative by clinical virologists). These are shown in Supplementary Table S3.

The R_t , which can be interpreted as the number of secondary infections caused by a person infected at a given date t , was estimated using national hospital admission data and the EpiEstim method [6,7]. Furthermore, the time series analysis to explore the added value of Cq data was performed using autoregressive integrated moving average (ARIMA) models. Further details about the methods can be found in the Supplementary Methods.

Ethical statement

This study was approved by the Institutional Review Board of the Centre Hospitalier Universitaire de Montpellier. It is registered at ClinicalTrials.gov under the identifier NCT04738331.

Results

Factors associated with Cq values

The adjusted R-square of the linear model was 38.8%, meaning that the factors we chose explained one-third of the variance in Cq values. The model residuals were normally distributed (Supplementary Figure S3A). Care should be taken in the interpretation given that the data were unbalanced, which is why we performed an analysis of variance (ANOVA) with type II sums of squares. All factors except the R_t were significantly associated with Cq values using a classical 5% p value criterion. Even for the control variable, the p value was 0.013, and patients with final digits 1 and 3 in their identification number had Cq values slightly lower (-0.19 and -0.17 cycles) than patients with a 0 as the final digit. Therefore, we set our significance thresholds to 5% of that of the control variables, i.e. 6.5×10^{-4} , to analyse the main effects (Table). Detailed outputs of the linear model are shown in Supplementary Table S2.

The intercept of the linear model indicates the average Cq value for a positive test performed with the reference assay, with all factors being set to their reference value. Its magnitude (19.1 cycles) is in line with clinical practice. The importance of the noise in the dataset is illustrated by the strong effect of the testing laboratory, as well as the RT-PCR assay used (Supplementary Figure S1 provides the distributions of Cq values as a function of the assay used and its target gene).

Despite this high level of noise, we detected a strong effect of the clinical qualitative result ('negative', 'positive' or 'weak positive') communicated by the testing laboratory (Figure 1A), with Cq differences that were even larger than those from the laboratory effect. We also found a significant difference of -1.81 cycles between the most common type of samples (nasopharyngeal) and other clinical sampling sites (mostly lower respiratory tract, but also faeces or saliva).

The effect associated with the number of days since symptom onset was particularly strong. The number of days between symptoms onset and testing dates was known for 8.5% of the participants; their Cq values increased gradually over the reported range of days with a maximum difference of 5.73 cycles (Figure 1B). The effect was similar when removing from the analysis the tests clinically considered to be 'negative' (see Supplementary Table S3 for these sensitivity analyses).

The effect of sex had the same order of magnitude as that of the control variable and could therefore be treated as non-significant. Conversely, the factor age

had a strong effect, with a decrease of 0.541 cycles per 20.1 years of age increased (Figure 1C).

The target gene of the RT-PCR assay used also yielded a significant effect. The Cq values obtained when using a probe targeting the ORF1 and S regions of the virus genome were significantly higher than when using the N gene, which was the genomic region of reference in the model (Figure 1D).

Finally, we found that Cq values decreased with time (-0.797 cycle per day), but this effect was nonlinear (Figure 1E). This could be due to the strong variation in testing efforts in France (Supplementary Figure S2A shows that the number of tests performed varied strongly during the year), but also to variations in the epidemic trend. Indeed, although the R_t (inferred from hospitalisation data) on the date of sampling was not significantly associated with the Cq value according to our threshold (6.5×10^{-4} , i.e. 5% of the lowest p value of the control variables), the interaction between the sampling date and R_t was nearly significant with a p value of 10^{-3} (Figure 1F), suggesting that a temporal analysis could yield additional insights.

Anticipating epidemic spread using Cq values

The existence of a correlation between the R_t and the Cq values of the tests performed in a population is consistent with population dynamics theory, which predicts that in an expanding population of infected individuals, the ‘age’ of the infections, i.e. the number of days post infection, is skewed towards lower values [8]. Since Cq values have been reported to increase over the course of an infection [3], which we confirm with this analysis (Figure 1D), it has been suggested that these values could be used as an early signal to predict R_t [9].

To investigate this question, we focused on screening data collected in the general population only from individuals aged 5 to 80 years because younger or older individuals may be associated with specific epidemiological clusters, e.g. in daycare facilities or nursing homes. We estimated the median and skewness values of the daily distribution of the Cq values. To correct for potential confounding factors, these were adjusted using a linear model (see details in the Supplementary Methods). We analysed the temporal correlation between the time series with a 7-day rolling average of this median, skew and R_t (Figure 2). For the median Cq value, we found a significant correlation with R_t that was maximised for a delay of 6–7 days (Supplementary Figure S4 provides additional information on the cross-correlation functions between R_t and the median or the skewness of the Cq distribution). This is consistent with R_t being calculated using data from hospital admissions for COVID-19, which occur at a median of 10 to 14 days after infection [10,11] and with RT-PCR screening data being obtained earlier in the infection. To further assess the usefulness of Cq data, we used ARIMA models to predict R_t dynamics over 7 days.

We compared models without any exogenous data to models that also included exogenous time series (either median or skewness of estimated Cq values distribution, or the fraction of positive tests [1]). As expected, the prediction error made using only endogenous data (R_t) was low in periods where R_t variations were limited. Furthermore, we found that adding exogenous data improved the prediction, especially when strong shifts in R_t were occurring (Figure 2). The Cq values tended to provide a better reduction in the error of the prediction than the ratio of positive tests.

Discussion

This analysis of a large database of RT-PCR tests performed in all of the French regions during the first two COVID-19 pandemic waves in 2020 confirms that population-level Cq values are noisy since even a linear model that featured 91 degrees of freedom did not explain the majority of the variance. However, owing to the law of large numbers, we detected several effects that are in line with biological observations and with virological properties. For instance, our finding that Cq values decreased as a function of the number of days after symptoms onset is consistent with longitudinal follow-up [3]. Another study also reported lower Cq values when the test was performed in symptomatic individuals [12]. The same study found that men had slightly lower Cq values than women, which was not significant in our analysis. Similarly, the difference we detected depending on the virus gene targeted by the RT-PCR assay used can be interpreted in the light of known differences in mRNA copy numbers between genes depending on their distance from the 3' end [13]. We also found slightly higher Cq values in samples collected from nasopharyngeal swabs compared with other samples (mainly lower respiratory tracts), but this is probably because the latter tests were performed in patients with more severe symptoms. Regarding the link we found between age and Cq values, although there are some mechanistic hypotheses as to why virus load would increase with age, such as variations in ACE2 receptor expression and immunosenescence [14], the evidence was mixed, with some studies reporting a decreasing trend [15] and others not [12,16]. Here, using a multivariate approach on a large dataset allowed us to unravel a strong and significant decrease of Cq values with age.

Finally, we found Cq differences associated with the gene targeted by the RT-PCR assay that are consistent with the life cycle of the virus. As stressed by Michalakis et al. [4], since coronaviruses are (+)ssRNA viruses, they use the same RNA matrix for replication and transcription, both being amplified by diagnostic assays. The problem is that the RNA matrix for transmission is not the same for each gene as *Coronaviridae* transcripts can produce subgenomic mRNAs that lack part of the genome [17]. As a consequence, and as shown in cell cultures [13], genes at the 5' end of the genome are under-represented. This is consistent with our result where assays targeting the gene at the 3' end (the N

gene) tend to have lower Cq values than assays targeting genes at the 5' end (the ORF1 and S genes). Note that an alternative explanation could be that some probes target more conserved areas of the SARS-CoV-2 than others [18].

A limitation of our study is that although our dataset stands out by its size and its level of details, it is restricted to a single country where testing effort varied, both on a temporal and on a spatial scale (Supplementary Figure S2). Although the testing behaviour can be assumed to be homogeneous, epidemic spread was different between regions, which could blur the relationship between Cq values and R_t at the national level. Performing similar analyses in other European countries and regions can also be particularly informative. In our study, we chose to analyse all the tests performed that had a Cq value. This is debatable since high Cq values can be due to noise and this is the point of implementing cut-offs. However, Cq values are known to increase during the course of an infection [3] and these high values could correspond to patients detected in a late stage, which is expected to be more frequent in a declining epidemic [9]. To control for this potential bias, we also performed the analysis on a dataset without the tests with a 'negative' result. Finally, this analysis was conducted at the end of the year 2020 but since then, as in most countries, the emergence of variants has altered epidemiological dynamics in France [19,20] and early reports suggest that the Cq value measured could depend on the variant causing the infection [21-25]. Vaccination has also changed the picture as indicated by Cq estimations in vaccine breakthrough infections [24,25] and should be included as a host factor in future analyses.

As pointed out elsewhere, care should be taken when interpreting Cq values because of technical issues (different assays may yield higher or lower values) and biological issues (coronaviruses produce subgenomic RNAs of different lengths) [4]. However, in this analysis, we do not attempt to link Cq values to viral loads but rather analyse raw values at a population level. A promising output of this analysis is the possibility to use Cq values as an early signal to detect changes in epidemic behaviour, e.g. R_t values. Indeed, in 2020, our most robust descriptors of the epidemic on a short time scale originated from hospital admission data, but these still have a considerable delay relative to the status of the epidemic since patients are hospitalised 2 weeks after infection [10,11]. The ratio of positive tests performed in the population of interest can, in theory, provide earlier insights but it suffers from strong sampling biases. We show that accounting for population-based variations in Cq values can improve R_t predictions on a 7-day period. This is consistent with a recent study which found a correlation between R_t and Cq distribution skewness using data from nasopharyngeal specimens collected from staff and residents in four long-term care facilities in Massachusetts, United States [9]. Note that, contrarily to us, this earlier study

did not factor in individual data such as patient age or symptomatic status, and it did not perform a cross-validation analysis that would control for temporal autocorrelation issues.

Our results show that analysing a large dataset of Cq values from screening tests can filter out the important amount of noise in these values. Inclusion of Cq values in routine surveillance calls for an adaptation to the current state of the epidemic, especially the evolution of variants and the increase in vaccination coverage, but also the integration with other types of data such as mobility data [26-28].

Conclusion

In many European Union countries, the qualitative outcome of SARS-CoV-2 screening tests are already aggregated in national databases to monitor epidemic spread. Adding Cq values as well as basic metadata (such as the RT-PCR assay used or, to a lesser extent, the age and sex) could be done while there should be minimal economical and ethical challenges. Our results call for a better integration of Cq values in national and European surveillance programmes to monitor epidemics caused by SARS-CoV-2 or other human viruses, especially since these data raise fewer ethical concerns than other sources such as mobility data.

Acknowledgements

We thank the ETE modelling team for discussion and the COVID-19 study group from the Société Française de Microbiologie (SFM) for their support.

Conflict of interest

None declared.

Authors' contributions

SA, VF, and SB conceived the study, SH-B, J-MG, LN, SP, TV, AdR, CT, MSo, RS, CB-B, MSa, A-SLH, SB, CL, JD, SH, CH, DV, HMD, SF, BV, CH, BL, VF, and SB contributed anonymous data, SA compiled the anonymous data sets from the collaborating partners, SA, MTS, and CS analysed the data, SA wrote the first draft of the manuscript, all authors commented and approved the manuscript.

References

1. Hasell J, Mathieu E, Beltekian D, Macdonald B, Giattino C, Ortiz-Ospina E, et al. A cross-country database of COVID-19 testing. *Sci Data*. 2020;7(1):345. <https://doi.org/10.1038/s41597-020-00688-8> PMID: 33033256
2. He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med*. 2020;26(5):672-5. <https://doi.org/10.1038/s41591-020-0869-5> PMID: 32296168
3. Néant N, Lingas G, Le Hingrat Q, Ghosn J, Engelmann I, Lepiller Q, et al. Modeling SARS-CoV-2 viral kinetics and association with mortality in hospitalized patients from the French COVID cohort. *Proc Natl Acad Sci USA*. 2021;118(8):e2017962118. <https://doi.org/10.1073/pnas.2017962118> PMID: 33536313
4. Michalakakis Y, Sofonea MT, Alizon S, Bravo IG. SARS-CoV-2 viral RNA levels are not 'viral load'. *Trends Microbiol*.

- 2021;29(11):970-2. <https://doi.org/10.1016/j.tim.2021.08.008> PMID: 34535373
5. French Microbiology Society (SFM). Avis du 25 septembre 2020 de la Société Française de Microbiologie (SFM) relatif à l'interprétation de la valeur de Ct (estimation de la charge virale) obtenue en cas de RT-PCR SARS-CoV-2 positive sur les prélèvements cliniques réalisés à des fins diagnostiques ou de dépistage. [Opinion on 25 September 2020 of the French Society of Microbiology (SFM) relating to the interpretation of the Ct value (estimate of the viral load) obtained in the event of a positive SARS-CoV-2 RT-PCR on clinical samples taken for diagnostic or screening purposes]. Paris: SFM; 2021. French. Available from: https://www.sfm-microbiologie.org/wp-content/uploads/2021/01/Avis-SFM-valeur-Ct-excre%CC%81tion-virale_-_Version-def-14012021_V4.pdf
 6. Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to estimate time-varying reproduction numbers during epidemics. *Am J Epidemiol.* 2013;178(9):1505-12. <https://doi.org/10.1093/aje/kwt133> PMID: 24043437
 7. Thompson RN, Stockwin JE, van Gaalen RD, Polonsky JA, Kamvar ZN, Demarsh PA, et al. Improved inference of time-varying reproduction numbers during infectious disease outbreaks. *Epidemics.* 2019;29:100356. <https://doi.org/10.1016/j.epidem.2019.100356> PMID: 31624039
 8. Caswell H. Matrix population models: construction, analysis and interpretation. Sunderland: Sinauer Associates, Inc.; 1989.
 9. Hay JA, Kennedy-Shaffer L, Kanjilal S, Lennon NJ, Gabriel SB, Lipsitch M, et al. Estimating epidemiologic dynamics from cross-sectional viral load distributions. *Science.* 2021;373(6552):eabho635. <https://doi.org/10.1126/science.abho635> PMID: 34083451
 10. Sofonea MT, Reyné B, Elie B, Djidjou-Demasse R, Selinger C, Michalakis Y, et al. Memory is key in capturing COVID-19 epidemiological dynamics. *Epidemics.* 2021;35:100459. <https://doi.org/10.1016/j.epidem.2021.100459> PMID: 34015676
 11. Salje H, Tran Kiem C, Lefrancq N, Courtejoie N, Bosetti P, Paireau J, et al. Estimating the burden of SARS-CoV-2 in France. *Science.* 2020;369(6500):208-11. <https://doi.org/10.1126/science.abc3517> PMID: 32404476
 12. Walker AS, Pritchard E, House T, Robotham JV, Birrell PJ, Bell I, et al. Ct threshold values, a proxy for viral load in community SARS-CoV-2 cases, demonstrate wide variation across populations and over time. *eLife.* 2021;10:e64683. <https://doi.org/10.7554/eLife.64683> PMID: 34250907
 13. Finkel Y, Mizrahi O, Nachshon A, Weingarten-Gabbay S, Morgenstern D, Yahalom-Ronen Y, et al. The coding capacity of SARS-CoV-2. *Nature.* 2021;589(7840):125-30. <https://doi.org/10.1038/s41586-020-2739-1> PMID: 32906143
 14. Yang Y, Zhao Y, Zhang F, Zhang L, Li L. COVID-19 in elderly adults: clinical features, molecular mechanisms, and proposed strategies. *Aging Dis.* 2020;11(6):1481-95. <https://doi.org/10.14336/AD.2020.0903> PMID: 33269102
 15. Euser S, Aronson S, Manders, I Lelyveld Sv, Herpers B, Sinnige J, et al. SARS-CoV-2 viral load distribution reveals that viral loads increase with age: a retrospective cross-sectional cohort study. *medRxiv.* 2021.01.15.21249691. preprint. <https://doi.org/10.1101/2021.01.15.21249691>
 16. Jones TC, Mühlemann B, Veith T, Biele G, Zuchowski M, Hofmann J, et al. An analysis of SARS-CoV-2 viral load by patient age. *medRxiv.* 2020.06.08.20125484. preprint. <https://doi.org/10.1101/2020.06.08.20125484>
 17. Ogando NS, Dalebout TJ, Zevenhoven-Dobbe JC, Limpens RWAL, van der Meer Y, Caly L, et al. SARS-coronavirus-2 replication in Vero E6 cells: replication kinetics, rapid adaptation and cytopathology. *J Gen Virol.* 2020;101(9):925-40. <https://doi.org/10.1099/jgv.0.001453> PMID: 32568027
 18. Dearlove B, Lewitus E, Bai H, Li Y, Reeves DB, Joyce MG, et al. A SARS-CoV-2 vaccine candidate would likely match all currently circulating variants. *Proc Natl Acad Sci USA.* 2020;117(38):23652-62. <https://doi.org/10.1073/pnas.2008281117> PMID: 32868447
 19. Haim-Boukobza S, Roquebert B, Trombert-Paolantoni S, Lecorche E, Verdurme L, Foulongne V, et al. Detecting Rapid Spread of SARS-CoV-2 Variants, France, January 26-February 16, 2021. *Emerg Infect Dis.* 2021;27(5):1496-9. <https://doi.org/10.3201/eid2705.210397> PMID: 33769253
 20. Alizon S, Haim-Boukobza S, Foulongne V, Verdurme L, Trombert-Paolantoni S, Lecorche E, et al. Rapid spread of the SARS-CoV-2 Delta variant in some French regions, June 2021. *Euro Surveill.* 2021;26(28):2100573. <https://doi.org/10.2807/1560-7917.ES.2021.26.28.2100573> PMID: 34269174
 21. Davies NG, Jarvis CI, Edmunds WJ, Jewell NP, Diaz-Ordaz K, Keogh RH, et al. Increased mortality in community-tested cases of SARS-CoV-2 lineage B.1.1.7. *Nature.* 2021;593(7858):270-4. <https://doi.org/10.1038/s41586-021-03426-1> PMID: 33723411
 22. Faria NR, Mellan TA, Whittaker C, Claro IM, Candido DDS, Mishra S, et al. Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus, Brazil. *Science.* 2021;372(6544):815-21. <https://doi.org/10.1126/science.abh2644> PMID: 33853970
 23. Roquebert B, Haim-Boukobza S, Trombert-Paolantoni S, Lecorche E, Verdurme L, Foulongne V, et al. SARS-CoV-2 variants of concern are associated with lower RT-PCR amplification cycles between January and March 2021 in France. *medRxiv.* 2021.03.19.21253971. preprint. <https://doi.org/10.1101/2021.03.19.21253971>
 24. Brown CM, Vostok J, Johnson H, Burns M, Gharpure R, Sami S, et al. Outbreak of SARS-CoV-2 infections, including COVID-19 vaccine breakthrough infections, associated with large public gatherings - Barnstable County, Massachusetts, July 2021. *MMWR Morb Mortal Wkly Rep.* 2021;70(31):1059-62. <https://doi.org/10.15585/mmwr.mm7031e2> PMID: 34351882
 25. Blanquart F, Abad C, Ambroise J, Bernard M, Cosentino G, Giannoli J-M, et al. Characterisation of vaccine breakthrough infections of SARS-CoV-2 Delta and Alpha variants and within-host viral load dynamics in the community, France, June to July 2021. *Euro Surveill.* 2021;26(37):2100824. <https://doi.org/10.2807/1560-7917.ES.2021.26.37.2100824> PMID: 34533119
 26. Kraemer MUG, Hill V, Ruis C, Dellicour S, Bajaj S, McCrone JT, et al. Spatiotemporal invasion dynamics of SARS-CoV-2 lineage B.1.1.7 emergence. *Science.* 2021;373(6557):889-95. <https://doi.org/10.1126/science.abj0113> PMID: 34301854
 27. Pullano G, Di Domenico L, Sabbatini CE, Valdano E, Turbelin C, Debin M, et al. Underdetection of cases of COVID-19 in France threatens epidemic control. *Nature.* 2021;590(7844):134-9. <https://doi.org/10.1038/s41586-020-03095-6> PMID: 33348340
 28. Selinger C, Choisy M, Alizon S. Predicting COVID-19 incidence in French hospitals using human contact network analytics. *Int J Infect Dis.* 2021;111:100-7. <https://doi.org/10.1016/j.ijid.2021.08.029> PMID: 34403783

License, supplementary material and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence and indicate if changes were made.

Any supplementary material referenced in the article can be found in the online version.

This article is copyright of the authors or their affiliated institutions, 2022.